

First time? Start here.

Before using your bottle for the first time, follow these steps to properly prepare.

Remove the cover and tab

Pull the clear cover straight up, then pull the green tab away from your bottle and discard both.



Uncap the green cap

Pull the cap straight up. Save it to protect your bottle for future use.



Prime

Dispense two drops onto a tissue. Flip your bottle upside down and slowly squeeze to ensure drops dispense one at a time.



To prevent contamination, do not let the tip of your bottle touch any surfaces.

Tips

- ✓ Turn your bottle upside down a few times to mix medicine before use
- ✓ Wait a few seconds after squeezing for drop to fully dispense
- ✓ Squeeze bottle slowly to dispense one drop at a time
- ✓ Let your bottle reinflate between drops and before recapping

**Congratulations, your multidose bottle is all set.
See reverse side for daily use instructions.**

APPROVED USE

RESTASIS MultiDose® Ophthalmic Emulsion helps increase your eyes' natural ability to produce tears, which may be reduced by inflammation due to Chronic Dry Eye. RESTASIS MultiDose® did not increase tear production in patients using anti-inflammatory eye drops or tear duct plugs.

IMPORTANT SAFETY INFORMATION

Do not use RESTASIS MultiDose® Ophthalmic Emulsion if you are allergic to any of the ingredients. Be careful not to touch the container tip to your eye or other surfaces, to help avoid eye injury and contamination.

RESTASIS MultiDose® should not be used while wearing contact lenses. If contact lenses are worn, they should be removed prior to use of RESTASIS MultiDose® and may be reinserted after 15 minutes.

PLEASE SEE ADDITIONAL IMPORTANT SAFETY INFORMATION ON BACK

Everyday use instructions

Your multidose bottle has a unique design.
For best results, follow these steps every time.

Uncap

Pull the cap straight up. Save it to protect your bottle for future use.



Drop

Dispense one drop in each eye. Flip your bottle upside down and slowly squeeze to ensure drops dispense one at a time.



Recap

Push the cap straight down, after your bottle reinflates to its original shape.



Get your Savings Card today



TEXT

Text SAVE to 72428*



ONLINE

Visit restasis.com/savings

IMPORTANT SAFETY INFORMATION (CONTINUED)

The most common side effect is a temporary burning sensation. Other side effects include eye redness, discharge, watery eyes, eye pain, foreign body sensation, itching, stinging, and blurred vision.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

PLEASE SEE ACCOMPANYING FULL PRODUCT INFORMATION

*By texting SAVE to 72428, you consent to receive recurring automated text messages from us and our partners. Your consent is not a condition of purchase of goods or services and you may opt out at any time. Message and data rates may apply. Message frequency depends on user. Mobile Terms and Conditions and Privacy Policy apply.